

HARMONIC MAPS AND SECTIONS ON SPHERES

M. BENYOUNES, E. LOUBEAU, AND C. M. WOOD

ABSTRACT. The absence of interesting harmonic sections for the Sasaki and Cheeger-Gromoll metrics has led to the consideration of alternatives, for example in the form of a two-parameter family of natural metrics shown to relax existence conditions for harmonicity [1].

This article investigates harmonic Killing vector fields, proves their non-existence on $\mathbb{S}^2$, obtains rigidity results for harmonic gradient vector fields on the two-sphere, classifies spherical quadratic gradient fields in all dimensions and determines the tension field, concluding with the discovery of a family of metrics making Hopf vector fields harmonic maps on $\mathbb{S}^{2n+1}$.

1. INTRODUCTION

Since the tangent bundle TM of a manifold M^n can be equipped with the structure of a $2n$ -dimensional manifold, it admits, at each of its points e , a tangent space of its own, denoted $T_e TM$, their disjoint union making up the bitangent bundle TTM .

For a Riemannian manifold (M, g) , the differential of the canonical projection $\pi : TM \rightarrow M$ and the Levi-Civita connection ∇ , via the connection map $K : TTM \rightarrow TM$, induce a natural splitting of TTM into vertical and horizontal subbundles so that:

$$T_e TM = V_e \oplus H_e.$$

As the restriction of K to V_e leads to a canonical identification with $T_x M$ ($\pi(e) = x$), we can lift vector fields on M to vertical and horizontal vector fields on V and H (cf. [4] for details).

Date: February 2, 2008.

1991 Mathematics Subject Classification. 58E20.

Key words and phrases. Harmonic sections, harmonic maps, Cheeger-Gromoll metrics.

The simplest metric one can define on TM is the Sasaki metric, which requires the projections from V and H to TM to be isometries and the two bundles to be orthogonal. Unfortunately, at least on compact manifolds, this most natural of metrics does not allow the existence of interesting harmonic maps or even sections (see definitions below). In order to obtain some flexibility, a new (two-parameter) family of metrics on TM was introduced in [1]:

$$h_{p,q}(A, B) = g(d\pi(A), d\pi(B)) + \omega^p(e) [g(KA, KB) + qg(KA, e)g(KB, e)],$$

where, $e \in TM$, $A, B \in T_e TM$, $\omega(e) = (1 + |e|^2)^{-1}$ and $p, q \in \mathbb{R}$.

While it is possible to allow q negative, $h_{p,q}$ being then only Riemannian in a tubular neighbourhood of the zero section, we will assume $q \geq 0$, except for a few special cases.

Note that $h_{0,0}$ is precisely the Sasaki metric and $h_{1,1}$ the Cheeger-Gromoll metric (cf. [6, 5] for more on their geometries).

These metrics $h_{p,q}$, named “generalised Cheeger-Gromoll metrics”, are all natural metrics (i.e. $d\pi$ is Riemannian and V and H are orthogonal). Their geometry is studied in [2].

2. (p, q) -HARMONIC SECTIONS

Assuming the base compact (otherwise work with compact subsets), we consider, rather than the “full” Dirichlet energy functional, the vertical energy of a vector field σ :

$$E^v(\sigma) = \frac{1}{2} \int_M |d^v \sigma|^2 v_g$$

(these two only differ by an additive constant) and derive its Euler-Lagrange equation for variations through sections [1]:

$$(1) \quad (1 + |\sigma|^2) \nabla^* \nabla \sigma + 2p \nabla_{X(\sigma)} \sigma = [p |\nabla \sigma|^2 - pq |X(\sigma)|^2 - q(1 + |\sigma|^2) \Delta(|\sigma|^2/2)] \sigma$$

where $\nabla^* \nabla \sigma = -\text{trace}(\nabla^2) \sigma$ (same sign convention for the Laplacian on functions) and $X(\sigma) = \nabla \left(\frac{|\sigma|^2}{2} \right) = g(\nabla \sigma, \sigma)$.

Clearly, parallel vector fields (when they exist) are (p, q) -harmonic sections and the first non-parallel examples are the standard Hopf vector

fields on $\mathbb{S}^{2p+1}$ ($p = 2$ and any q). Besides, rescaling them yields examples for all (p, q) with $p > 1$. Moreover, violating our convention $q \geq 0$, conformal vector fields on $\mathbb{S}^n$ are (p, q) -harmonic if and only if $n \geq 3$, $p = n + 1$ and $q = 2 - n$ ([1]).

On the other hand, when the base is compact (without boundary) and $p \leq 1$ ($q \geq 0$), then a vector field is (p, q) -harmonic if and only if it is parallel (cf. [1]).

This includes the case of the original Cheeger-Gromoll metric ($p = q = 1$) (see also [7]). Note our convention for the curvature tensor: $R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}$.

3. KILLING VECTOR FIELDS

As illustrated by Hopf fields on $\mathbb{S}^{2p+1}$, Killing vector fields are prime candidates for (p, q) -harmonicity.

Proposition 3.1. *A Killing vector field σ is (p, q) -harmonic if and only if:*

$$(2) \quad (1 + |\sigma|^2) \text{Ricci}(\sigma) - 2p \nabla_{\nabla_\sigma \sigma} \sigma = \left[p |\nabla \sigma|^2 - pq |\nabla_\sigma \sigma|^2 - q(1 + |\sigma|^2) \Delta(\tfrac{1}{2} |\sigma|^2) \right] \sigma$$

Proof. For σ Killing, $X(\sigma) = -\nabla_\sigma \sigma$ and $\nabla_{X, Y}^2 \sigma = -R(\sigma, X)Y$ so $\nabla^* \nabla \sigma = \text{Ricci}(\sigma)$. $\square$

The first sub-case to investigate is constant norm.

Proposition 3.2. *A Killing vector field σ of constant norm $|\sigma|^2 = k^2$ is (p, q) -harmonic if and only if it is an eigenvector of the Ricci operator. Then $p = 1 + \frac{1}{k^2}$ and q is any (positive) real number.*

Proof. If σ is Killing and $|\sigma|^2 = k^2$, then $X(\sigma) = 0$ and $\text{Ricci}(\sigma, \sigma) = |\nabla \sigma|^2$ ([8]), so (p, q) -harmonicity becomes:

$$(1 + k^2) \text{Ricci}(\sigma) = (p |\nabla \sigma|^2) \sigma = p \text{Ricci}(\sigma, \sigma) \sigma.$$

$\square$

Example 3.1. There are easy examples of Killing vector fields which are eigenvectors of Ricci on the space forms $\mathbb{R}^n$ and $\mathbb{S}^{2p+1}$. In dimension three, one can also search among Thurston's geometries, starting with

$\mathbb{S}^2 \times \mathbb{R}$ and $\frac{\partial}{\partial t}$. Unfortunately, all these examples, except Hopf fields, are parallel.

To obtain new non-parallel examples, we turn to the Heisenberg space H_3 , identified with $\mathbb{R}^3$ endowed with the metric $dx^2 + dy^2 + (dz - xdy)^2$ and $\widetilde{SL_2(\mathbb{R})}$, the universal cover of the special linear group, seen as $\mathbb{R}_+^3 = \{(x, y, z) \in \mathbb{R}^3 : z > 0\}$ with the metric $(dx + \frac{dy}{z})^2 + \frac{dy^2 + dz^2}{z^2}$. For H_3 , the vectors $E_1 = \frac{\partial}{\partial x}$, $E_2 = \frac{\partial}{\partial y} + x\frac{\partial}{\partial z}$, $E_3 = \frac{\partial}{\partial z}$ form an orthonormal basis and E_3 is Killing and non-parallel. Besides one verifies that $\text{Ricci}(E_3) = \frac{1}{2}E_3$.

For $\widetilde{SL_2(\mathbb{R})}$, things are similar, since an orthonormal basis is given by $E_1 = \frac{\partial}{\partial x}$, $E_2 = z\frac{\partial}{\partial y} - \frac{\partial}{\partial x}$, $E_3 = z\frac{\partial}{\partial z}$ and E_1 is Killing and non-parallel, moreover $\text{Ricci}(E_1) = \frac{1}{2}E_1$.

More generally, constancy of the norm is replaced with harmonicity.

Proposition 3.3. *Let σ be a (p, q) -harmonic Killing vector field with harmonic norm. If $p \geq 0$ and $(p - 1)|\sigma|^2 \leq 1$ (automatic when $p \leq 1$) then σ is of constant norm.*

Proof. In this case, $\text{Ricci}(\sigma, \sigma) = |\nabla \sigma|^2$ and σ (p, q) -harmonic implies:

$$(1 + (1 - p)|\sigma|^2) \text{Ricci}(\sigma, \sigma) = -p(2 + q|\sigma|^2)|\nabla \sigma|^2$$

hence σ has constant norm. $\square$

Remark 3.1. On compact manifolds, Proposition 3.3 is only a minute improvement on [1, Theorem 4.6].

Remark 3.2. Non-parallel Killing vector fields do not exist on a compact manifold with negative Ricci. On the other hand, when $\text{Ricci} > 0$, if $|\sigma|^2$ is harmonic then it must be constant [9].

The two-sphere is a particularly restrictive case.

Proposition 3.4. *There is no non-zero (p, q) -harmonic Killing vector field on $\mathbb{S}^2$ ($p > 0$, $q \geq 0$).*

We first need:

Lemma 3.1. *Let σ be a non-zero Killing vector field on $\mathbb{S}^2$. Then:*

- i) *The set $A = \{x \in \mathbb{S}^2 : \sigma(x) = 0\}$ has empty interior.*
- ii) *If $\nabla_\sigma \sigma = 0$ on an open subset U then $\Delta \frac{|\sigma|^2}{2} = 0$ on U .*

Proof. i) If $\sigma \equiv 0$ on an open subset B of $\mathbb{S}^2$, then $(\nabla_X \sigma)(x) = 0, \forall x \in B, \forall X \in T_x \mathbb{S}^2$ and $\sigma \equiv 0$ on $\mathbb{S}^2$ ([8]).

ii) Assume $\nabla_\sigma \sigma \equiv 0$ on an open subset U of $\mathbb{S}^2$, then $|\sigma|^2$ is constant and $\Delta \frac{|\sigma|^2}{2} = 0$ on U . $\square$

Proof of Proposition 3.4. Let σ be a non-zero (p, q) -harmonic Killing vector field on $\mathbb{S}^2$, U and A as in the above lemma.

On $W = \mathbb{S}^2 \setminus (U \cup A)$, $\sigma/|\sigma|$ and $\nabla_\sigma \sigma/|\nabla_\sigma \sigma|$ form an orthonormal basis, put $V = \nabla_\sigma \sigma$ then:

$$|\nabla \sigma|^2 = \frac{1}{|\sigma|^2} |\nabla_\sigma \sigma|^2 + \frac{1}{|V|^2} |\nabla_V \sigma|^2.$$

Since σ is (p, q) -harmonic:

$$\nabla_{\nabla_\sigma \sigma} \sigma = \frac{-1}{2p} \left(p |\nabla \sigma|^2 - pq |\nabla_\sigma \sigma|^2 - q(1 + |\sigma|^2) \Delta \left(\frac{1}{2} |\sigma|^2 \right) - (1 + |\sigma|^2) \right) \sigma$$

so that $\nabla_{\nabla_\sigma \sigma} \sigma = f \sigma$ with $f = \frac{-1}{2p} \left(p |\nabla \sigma|^2 - pq |\nabla_\sigma \sigma|^2 - q(1 + |\sigma|^2) \Delta \left(\frac{1}{2} |\sigma|^2 \right) - (1 + |\sigma|^2) \right)$. On W :

$$|\nabla_\sigma \sigma|^2 = -f |\sigma|^2 \quad \text{and} \quad |\nabla_V \sigma|^2 = f^2 |\sigma|^2.$$

Therefore $|\nabla \sigma|^2 = -2f$. Note that by continuity this holds on $\mathbb{S}^2 \setminus U$ and f cannot vanish.

Hence

$$0 = pq |\nabla_\sigma \sigma|^2 + q(1 + |\sigma|^2) \Delta \left(\frac{1}{2} |\sigma|^2 \right) + (1 + |\sigma|^2)$$

so that $|\sigma|^2$ is constant. $\square$

However, one can try to obtain (p, q) -harmonic vector fields by multiplying a Killing vector field by a function.

Example 3.2. Consider $\sigma(x, y, z) = (-y, x, 0)$ the Killing vector field on $\mathbb{S}^2$ given by rotation around the z -axis and complete it into an orthonormal frame $\left\{ \frac{\sigma}{x^2+y^2}, \frac{\gamma}{x^2+y^2} \right\}$ of $\mathbb{S}^2 \setminus \{N, S\}$, the sphere less the poles.

For $f \in C^\infty(\mathbb{S}^2)$, let $V = f \sigma$, then standard computations show that V (p, q) -harmonic requires $\sigma(f) = 0$, i.e. $f(x, y, z) = F(x^2 + y^2)$ for

some real function F .

Then $V = F(x^2 + y^2)\sigma$ is $(p, 0)$ -harmonic if:

$$(3) \quad (1+tF^2)[4t(t-1)F'' + (10t-8)F' + F] + 4p(1-t)[tF^2F' + t^2F(F')^2] = 0$$

Clearly constant functions are not solutions and while $F(t) = \frac{1}{\sqrt{(p-1)t}}$ is a solution, it does not extend over the poles.

More generally, if F is a solution of (3) on $[0, 1]$, assuming (wlog) $F(0) \geq 0$ and $F'(t) \geq 0$ for $t \in]0, \epsilon[$ ($\epsilon > 0$), then $F' \geq \frac{1}{\sqrt{t^4(1-t)}}$ on $]0, \epsilon[$. Hence no function F on $\mathbb{S}^2$ can make $F\sigma$ $(p, 0)$ -harmonic.

4. GRADIENT VECTOR FIELDS

First, we find an obstruction for another important class of vector fields on $\mathbb{S}^2$.

Proposition 4.1. *Let σ be a (non-zero) (p, q) -harmonic vector field defined by $\sigma = \nabla f$ with $\Delta f = \lambda f$. If $\nabla_\sigma \sigma$ and σ are non-colinear on an open subset U of $\mathbb{S}^2$, then $\Delta \frac{|\sigma|^2}{2} \leq 0$ on U .*

Proof. From standard properties of gradient vector fields, $\sigma = \nabla f$ with $\Delta f = \lambda f$, is (p, q) -harmonic if:

$$(1+|\sigma|^2)(\lambda-1)\sigma + 2p\nabla_{\nabla_\sigma \sigma}\sigma = \left[p|\nabla\sigma|^2 - pq|\nabla_\sigma \sigma|^2 - q(1+|\sigma|^2)\Delta \frac{|\sigma|^2}{2} \right] \sigma$$

Therefore $\nabla_{\nabla_\sigma \sigma}\sigma = f\sigma$ with:

$$f = \frac{1}{2p} \left(p|\nabla\sigma|^2 - pq|\nabla_\sigma \sigma|^2 - q(1+|\sigma|^2)\Delta \frac{|\sigma|^2}{2} - (\lambda-1)(1+|\sigma|^2) \right).$$

Note that σ cannot vanish on an open subset of $\mathbb{S}^2$. Assume that $\left\{ \frac{\sigma}{|\sigma|}, \frac{b}{|b|} \right\}$, where $b = \nabla_\sigma \sigma - \langle \nabla_\sigma \sigma, \sigma \rangle \frac{\sigma}{|\sigma|^2}$, is an orthonormal basis on an open subset U , then:

$$|\nabla\sigma|^2 = \left| \nabla \frac{\sigma}{|\sigma|} \sigma \right|^2 + \left| \nabla \frac{b}{|b|} \sigma \right|^2$$

with

$$\left| \nabla \frac{\sigma}{|\sigma|} \sigma \right|^2 = \frac{1}{|\sigma|^2} \langle f\sigma, \sigma \rangle = f$$

while

$$\left| \nabla \frac{b}{|b|} \sigma \right|^2 = \frac{1}{|b|^2} |\nabla_b \sigma|^2$$

$$|b|^2 = f|\sigma|^2 - \frac{1}{|\sigma|^2} \langle \nabla_\sigma \sigma, \sigma \rangle^2 \quad \text{and} \quad |\nabla_b \sigma|^2 = f \left[f|\sigma|^2 - \frac{1}{|\sigma|^2} \langle \nabla_\sigma \sigma, \sigma \rangle^2 \right]$$

Therefore $\frac{1}{|b|^2} |\nabla_b \sigma|^2 = f$ and $|\nabla \sigma|^2 = 2f$.

Plugging in the definition of f , we have:

$$0 = -pq|\nabla_\sigma \sigma|^2 - q(1 + |\sigma|^2) \Delta \frac{|\sigma|^2}{2} - (\lambda - 1)(1 + |\sigma|^2),$$

therefore $\Delta \frac{|\sigma|^2}{2} \leq 0$ on U .

If U is dense in $\mathbb{S}^2$, this forces $|\sigma|^2$ to be constant. $\square$

Remark 4.1. For conformal vector fields on $\mathbb{S}^n$, $\nabla_\sigma \sigma$ and σ are colinear. However [1, Theorem 5.4] states that conformal vector fields on $\mathbb{S}^2$ cannot be (p, q) -harmonic.

4.1. Spherical Quadratic Gradient Fields. To a symmetric bilinear form β on $\mathbb{R}^{n+1}$, associate the linear map B :

$$B(x) \cdot y = \beta(x, y), \quad \text{for all } x, y \in \mathbb{R}^{n+1},$$

define the restriction of the associated quadratic form to $\mathbb{S}^n$:

$$\lambda(x) = \beta(x, x) = Bx \cdot x,$$

and $\sigma = \frac{1}{2} \nabla \lambda$ its spherical gradient. Clearly $\sigma(x) = B(x) - \lambda(x)x$. More generally, for any $k \in \mathbb{N}$:

$$\lambda_k(x) = B^k(x) \cdot x, \quad \text{and} \quad \sigma_k(x) = \frac{1}{2} \nabla \lambda_k = B^k(x) - \lambda_k(x)x.$$

Consider the symmetric $(1, 1)$ -tensors on $\mathbb{S}^n$:

$$L_k X = B^k X - (B^k X \cdot x)x,$$

and put $L = L_1$. Note that: $L_k \neq L^k$ if $k > 1$.

By direct calculations, we obtain:

Proposition 4.2.

- (a) $\langle \sigma_k, \sigma_l \rangle = \lambda_{k+l} - \lambda_k \lambda_l$
- (b) $L_l \sigma_k = \sigma_{k+l} - \lambda_k \sigma_l$
- (c) $\nabla_X \sigma_k = L_k X - \lambda_k X$
- (d) $\operatorname{div} \sigma_k = \operatorname{trace}(B^k) - (n+1)\lambda_k$
- (e) $|L_k|^2 = |B^k|^2 - 2\lambda_{2k} + \lambda_k^2$
- (f) $\nabla_X L_k(Y) = -\langle X, Y \rangle \sigma_k - \langle \sigma_k, Y \rangle X$
- (g) $\nabla_{X,Y}^2 \sigma_k = -\langle X, Y \rangle \sigma_k - \langle \sigma_k, Y \rangle X - 2\langle \sigma_k, X \rangle Y$

Corollary 4.1. (a) When L_k is viewed as a TM -valued 1-form on

$$M: \quad \delta L_k = (n+1)\sigma_k$$

$$(b) \quad \nabla^* \nabla \sigma_k = (n+3)\sigma_k$$

A straightforward consequence of Proposition 4.2 is:

Proposition 4.3. For $\sigma = \frac{1}{2}\nabla\lambda$, we have:

- (a) $|\sigma|^2 = \lambda_2 - \lambda^2$
- (b) $X(\sigma) = \sigma_2 - 2\lambda\sigma$
- (c) $\nabla_X \sigma = LX - \lambda X$
- (d) $\nabla_{X(\sigma)} \sigma = \sigma_3 - 3\lambda\sigma_2 + (4\lambda^2 - \lambda_2)\sigma$
- (e) $|\nabla\sigma|^2 = |B|^2 - 2\lambda_2 - 2\lambda \operatorname{trace} B + (n+3)\lambda^2$
- (f) $\nabla^* \nabla \sigma = (n+3)\sigma$
- (g) $\Delta|\sigma|^2 = -2|B|^2 + 2(n+5)\lambda_2 + 4\lambda \operatorname{trace} B - 4(n+3)\lambda^2$

This formalism leads to a classification.

Theorem 4.1. A quadratic gradient vector field σ on $\mathbb{S}^n$ is (p, q) -harmonic if and only if n is odd, $n \geq 5$, $p = \frac{n+3}{2}$ and q is the only solution (necessarily negative) of:

$$8(n+1)q^2 + 2(3n^2 - 2n - 17)q + (n-3)(n^2 - 9) = 0$$

such that $\mu^2 = \frac{9-n^2}{4q} - 2(n+3)$ is strictly positive.

Besides, σ is given by $\frac{1}{2}\nabla\lambda$ where $\lambda(x) = Bx \cdot x$ and B is a symmetric matrix with two eigenvalues α and β , of equal multiplicities $\frac{n+1}{2}$, such that $|\alpha - \beta| = \mu$.

Corollary 4.2. *There is no (p, q) -harmonic quadratic gradient vector fields on even-dimensional spheres, and, more generally, none for positive values of q (recall that $h_{p,q}$ is a Riemannian metric on the whole of $T\mathbb{S}^n$ if and only if q is positive).*

Remark 4.2. a) B and $\tilde{B} = B - \gamma Id$ give rise to the same gradient vector field.

b) Since B can be written $B = P^{-1}AP$, A diagonal and P orthogonal, $\lambda(x) = B(x).x$ and $\tilde{\lambda}(x) = A(x).x$ are isometric and so are $\sigma = \frac{1}{2}\nabla\lambda$ and $\tilde{\sigma} = \frac{1}{2}\nabla\tilde{\lambda}$. More prosaically:

$$\lambda(x) = \langle B(x), x \rangle = \langle P^{-1}APx, x \rangle = \langle Ax, x \rangle \circ \phi,$$

for some isometry ϕ .

So, up to isometries and constants, the (p, q) -harmonic vector fields of Theorem 4.1 are the gradients of $\lambda(x) = x_1^2 + \dots + x_{\frac{n+1}{2}}^2$.

Proof of Theorem 4.1. From Proposition 4.3:

$$(4) \quad (1 + |\sigma|^2)\nabla^*\nabla\sigma + 2p\nabla_{X(\sigma)}\sigma = 2p\sigma_3 - 6p\lambda\sigma_2 + (n + 3 + (n + 3 - 2p)\lambda_2 + (8p - n - 3)\lambda^2)\sigma$$

hence, if σ is (p, q) -harmonic, $\sigma_3 - 3\lambda\sigma_2$ must be colinear to σ .

Remark 4.3. a) It is easy to see that $\sigma_2 = m\sigma$ for $m \in \mathbb{R}$ if and only if B satisfies $B^2 - mB = cI$ for $c \in \mathbb{R}$.

b) $B^2 - mB = cI$ is equivalent to B having either one or two distinct eigenvalues, depending on whether the roots of the polynomial $x^2 - mx - c$ are repeated or distinct. If B has a single eigenvalue μ then $m = 0$ and $c = \mu^2$, whereas if B has distinct eigenvalues μ, ν then $m = \mu + \nu$ and $c = -\mu\nu$, and:

$$\begin{aligned} \sigma(x) &= (\mu - \nu)|x_\nu|^2 x_\mu + (\nu - \mu)|x_\mu|^2 x_\nu \\ \sigma_2(x) &= (\mu^2 - \nu^2)|x_\nu|^2 x_\mu + (\nu^2 - \mu^2)|x_\mu|^2 x_\nu, \end{aligned}$$

where $x = x_\mu + x_\nu$ is the decomposition into eigenvectors.

Lemma 4.1. *A non-trivial σ satisfies:*

$$(5) \quad \sigma_3 - 3\lambda\sigma_2 = f\sigma,$$

for some smooth function $f: \mathbb{S}^n \rightarrow \mathbb{R}$ if and only if B has precisely two distinct eigenvalues.

Proof. If B has two distinct eigenvalues μ, ν , then $B^2 - mB = cI$, where $m = \mu + \nu$ and $c = -\mu\nu$ and:

$$\sigma_3 - 3\lambda\sigma_2 = (m^2 - 3m\lambda + c)\sigma$$

To show the converse, let $\{e_i\}$ be an orthonormal B -eigenbasis of $\mathbb{R}^{n+1}$:

$$Be_i = \mu_i e_i, \quad \text{where } \mu_i \in \mathbb{R}.$$

Notice that for all $k \in \mathbb{N}$ we have:

$$\lambda_k(e_i) = \mu_i^k \quad \text{and } \sigma_k(e_i) = 0,$$

then $x = e_i \in \mathbb{S}^n$ and covariantly differentiating Equation (5) along $X = e_j \in T_x \mathbb{S}^n$ for some $j \neq i$ yields:

$$\nabla_X \sigma_3 - 3\mu_i \nabla_X \sigma_2 = f(e_i) \nabla_X \sigma$$

which reduces to:

$$\mu_j^3 - \mu_i^3 - 3\mu_i(\mu_j^2 - \mu_i^2) = (\mu_j - \mu_i)f(e_i), \quad \text{for all } i \neq j.$$

If σ is non-trivial, there exist i, j such that $\mu_i \neq \mu_j$, and:

$$(6) \quad f(e_i) = \mu_j^2 - 2\mu_i\mu_j - 2\mu_i^2$$

$$(7) \quad f(e_j) = \mu_i^2 - 2\mu_i\mu_j - 2\mu_j^2$$

If μ_k is a third distinct eigenvalue then:

$$f(e_i) = \mu_k^2 - 2\mu_i\mu_k - 2\mu_i^2$$

and comparing with (6) and (7) yields:

$$\mu_i = \frac{1}{2}(\mu_j + \mu_k) \quad \text{and } \mu_j = \frac{1}{2}(\mu_i + \mu_k).$$

Therefore:

$$\mu_j = \frac{1}{2}(\mu_i + \mu_k) = \frac{1}{4}(\mu_j + 3\mu_k)$$

hence $\mu_j = \mu_k$. So B has at most two distinct eigenvalues. $\square$

End of proof of Theorem 4.1:

By Remark 4.2 a), one eigenvalue, ν , can be chosen zero, and the other, μ , strictly positive, of multiplicity k . Decomposing $x = x_\mu + x_\nu \in \mathbb{S}^n$ into eigenvectors, the left-hand side of (4) becomes:

$$(8) \quad 2p\mu^2 + n + 3 + (n + 3 - 8p)\mu^2|x_\mu|^2 + (8p - n - 3)\mu^2|x_\mu|^4$$

while its right-hand side gives:

$$(9) \quad \begin{aligned} & k\mu^2(p + q) + \mu^2[-2p(k + 1) - pq\mu^2 + kq\mu^2 - q(n + 5 + 2k)]|x_\mu|^2 + \\ & \mu^2[p(n + 3) + 5pq\mu^2 - q(n + 5 + 2k)\mu^2 - kq\mu^2 + 2(n + 3)q]|x_\mu|^4 + \\ & q\mu^4[-8p + 2(n + 3) + n + 5 + 2k]|x_\mu|^6 + 2q\mu^4(2p - n - 3)|x_\mu|^8 \end{aligned}$$

Note that by continuity, (8) and (9) must be equal everywhere. Equating the coefficients yields $p = \frac{n+3}{2}$, $k = \frac{n+1}{2}$ (but rules out $q = 0$) and the system:

$$(10) \quad \begin{cases} \frac{n^2-9}{4} + q(\mu^2 + 2(n + 3)) = 0 & (\text{forcing } q < 0 \text{ and excluding } n = 3) \\ 8(n + 1)q^2 + 2(3n^2 - 2n - 17)q + (n - 3)(n^2 - 9) = 0 \end{cases}$$

of the form:

$$(11) \quad \begin{cases} (\mu^2 + a)q = K \\ \mu^2(q + b) = K' \end{cases}$$

with K negative and K' positive. These represent two (branches) of hyperbolas in different quadrants but intersecting (once and only once).

□

5. HARMONIC MAPS

With the same energy functional E^v on sections, one can formulate several variational problems, according to the type of variations allowed. The lifting of all restrictions, that is permitting variations through mere maps, brings us back to harmonic maps (cf. [3]). In this situation, the associated Euler-Lagrange operator is the tension field $\tau(\sigma) = \text{trace } \nabla d\sigma$.

Standard computations provide:

Proposition 5.1. [2] *The Levi-Civita connection $\tilde{\nabla}$ of the generalised Cheeger-Gromoll metric $h_{p,q}$ is given by:*

$$\begin{aligned}\tilde{\nabla}_{X^h} Y^h &= (\nabla_X Y)^h - \frac{1}{2}(R(X, Y)e)^v \\ \tilde{\nabla}_{X^h} Y^v &= (\nabla_X Y)^v + \frac{\omega^p(e)}{2}(R(e, Y)X)^h \\ \tilde{\nabla}_{X^v} Y^h &= \frac{\omega^p(e)}{2}(R(e, X)Y)^h \\ \tilde{\nabla}_{X^v} Y^v &= -p\omega(e)[g(X, e)Y^v + g(Y, e)X^v] + \\ &\quad \frac{p\omega(e)+q}{1+q|e|^2}g(X, Y)U + \frac{pq\omega(e)}{1+q|e|^2}g(X, e)g(Y, e)U,\end{aligned}$$

where R is the curvature tensor of (M, g) and U the canonical vertical vector field.

Note that $\tilde{\nabla}_{X^v} Y^v \in V$ so the fibres are totally geodesic.

This enables us to compute the tension field:

Proposition 5.2. *A vector field $\sigma : (M, g) \rightarrow (TM, h_{p,q})$ is a harmonic map if and only if:*

$$(12) \quad \begin{cases} \sum_{i=1}^n R(\sigma, \nabla_{e_i} \sigma) e_i = 0 \\ \nabla^* \nabla \sigma + 2p\omega(\sigma) \nabla_{X(\sigma)} \sigma = \left[\frac{p\omega(\sigma)+q}{1+q|\sigma|^2} |\nabla \sigma|^2 + \frac{pq\omega(\sigma)}{1+q|\sigma|^2} |X(\sigma)|^2 \right] \sigma \end{cases}$$

where $\{e_i\}_{i=1, \dots, n}$ is an orthonormal frame on (M, g) .

Proof. A vector field being an immersive bijection, $\sigma(M) \subset TM$ is a submanifold of TM .

Therefore, using a orthonormal frame $\{e_i\}_{i=1, \dots, n}$ on M :

$$\begin{aligned}\tau(\sigma) &= \text{trace } \nabla d\sigma = \sum_{i=1}^n \tilde{\nabla}_{d\sigma(e_i)} d\sigma(e_i) - d\sigma(\nabla_{e_i} e_i) \\ &= \sum_{i=1}^n \tilde{\nabla}_{(e_i)^h + (\nabla_{e_i} \sigma)^v} [(e_i)^h + (\nabla_{e_i} \sigma)^v] - (\nabla_{e_i} e_i)^h - (\nabla_{\nabla_{e_i} e_i} \sigma)^v \\ &= \left[\nabla^* \nabla \sigma - 2p\omega(\sigma) \nabla_{X(\sigma)} \sigma + \frac{p\omega(\sigma)+q}{1+q|\sigma|^2} |\nabla \sigma|^2 \sigma + \frac{pq\omega(\sigma)}{1+q|\sigma|^2} |X(\sigma)|^2 \sigma \right]^v \\ &\quad + \sum_{i=1}^n \omega^m(e) (R(\sigma, \nabla_{e_i} \sigma) e_i)^h\end{aligned}$$

□

Since the vector tangent to a variation through sections must be vertical, the Euler-Lagrange equation of the functional E^v obtained by variations through sections (Equation (1)) must be equivalent to the vertical part of the tension field. The next proposition checks this directly.

Proposition 5.3. *A section σ of TM satisfies:*

(13)

$$(1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma=\left[p|\nabla\sigma|^2-pq|X(\sigma)|^2-q(1+|\sigma|^2)\Delta\frac{|\sigma|^2}{2}\right]\sigma$$

if and only if

$$(14) \quad \nabla^*\nabla\sigma+2p\omega(\sigma)\nabla_{X(\sigma)}\sigma=\left[\frac{p\omega(\sigma)+q}{1+q|\sigma|^2}|\nabla\sigma|^2+\frac{pq\omega(\sigma)}{1+q|\sigma|^2}|X(\sigma)|^2\right]\sigma$$

Proof. Assume (13) and $q \neq 0$, then (14) is equivalent to

$$\begin{aligned} & -q|\sigma|^2(1+|\sigma|^2)\nabla^*\nabla\sigma-2pq|\sigma|^2\nabla_{X(\sigma)}\sigma+q(1+|\sigma|^2)(\Delta\frac{|\sigma|^2}{2})\sigma \\ & +\left[q(1+|\sigma|^2)|\nabla\sigma|^2+2pq|X(\sigma)|^2\right]\sigma=0 \end{aligned}$$

$$\Leftrightarrow |\sigma|^2\left((1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma\right)=\left[(1+|\sigma|^2)(|\nabla\sigma|^2+\Delta\frac{|\sigma|^2}{2})+2p|X(\sigma)|^2\right]\sigma$$

$$\Leftrightarrow |\sigma|^2\left((1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma\right)=g((1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma,\sigma)\sigma$$

$$\text{since } g(\nabla_{X(\sigma)}\sigma,\sigma)=|X(\sigma)|^2$$

Therefore (14) is equivalent to $|\sigma|^2\Lambda=g(\Lambda,\sigma)\sigma$ where $\Lambda=(1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma$, and this is true if and only if $\Lambda=f\sigma$, which is precisely what (13) says.

Conversely, assume that (14) holds, then (13) is equivalent to

$$(1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma=$$

$$\left[p|\nabla\sigma|^2-pq|X(\sigma)|^2-q(1+|\sigma|^2)(g(\nabla^*\nabla\sigma,\sigma)-|\nabla\sigma|^2)\right]\sigma$$

$$\Leftrightarrow (1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma+q\left[g((1+|\sigma|^2)\nabla^*\nabla\sigma+p\nabla_{X(\sigma)}\sigma,\sigma)\right]\sigma$$

$$=(p+q+q|\sigma|^2)|\nabla\sigma|^2\sigma$$

$$\Leftrightarrow [(1+|\sigma|^2)\nabla^*\nabla\sigma+2p\nabla_{X(\sigma)}\sigma](1+q|\sigma|^2)=$$

$$[(p+q+q|\sigma|^2)|\nabla\sigma|^2+pq|X(\sigma)|^2]\sigma$$

since, again, (14) implies that $|\sigma|^2\Lambda=g(\Lambda,\sigma)\sigma$. □

Proposition 5.4. *The scaled Hopf vector fields are harmonic maps from $\mathbb{S}^{2n+1}$ to $T\mathbb{S}^{2n+1}$ equipped with the generalised Cheeger-Gromoll metric $h_{p,q}$ ($p > 1$, $q \geq 0$).*

Proof. On spheres, the horizontal part of the tension field becomes:

$$\sum_{i=1}^n R(\sigma, \nabla_{e_i} \sigma) e_i = (\operatorname{div} \sigma) \sigma - \nabla_{\sigma} \sigma$$

So if σ is Killing of constant norm, it is harmonic. $\square$

Remark 5.1. a) For $2 \geq p > 1$ and $q = 0$, these represent new examples of harmonic maps from a compact manifold into positive sectional curvature (cf. [2]).

b) No Killing vector field on $\mathbb{S}^{2n}$ can be a harmonic map.

Example 5.1. Proposition 5.4 also extends to the vector fields on Heisenberg and $\widetilde{SL_2(\mathbb{R})}$ considered in Example 3.1:

- i) H_3 : With the same notations, let σ be the vector E_3 , since the non-zero components of the curvature tensor of H_3 are $R_{1221} = \frac{3}{4}$ and $R_{1331} = -\frac{1}{4} = R_{2332}$, then $\sum_{i=1}^3 R(\sigma, \nabla_{e_i} \sigma) e_i = 0$.
- ii) $\widetilde{SL_2(\mathbb{R})}$: Still with the same notations as in Example 3.1, let σ be the vector E_1 , then: $\sum_{i=1}^3 R(\sigma, \nabla_{e_i} \sigma) e_i = 0$, since $R(E_1, E_3)E_2 = R(E_1, E_2)E_3 = 0$.

REFERENCES

- [1] M. Benyounes, E. Loubeau and C. M. Wood, Harmonic sections of Riemannian vector bundles and metrics of Cheeger-Gromoll type, to appear in *Diff. Geom. Appl.*
- [2] M. Benyounes, E. Loubeau and C. M. Wood, The geometry of generalised Cheeger-Gromoll metrics, in preparation.
- [3] J. Eells and L. Lemaire, *Selected topics on harmonic maps*, AMS, 1983.
- [4] S. Gudmundsson and E. Kappos, On the geometry of tangent bundles, *Expositiones Mathematicae*, 20 (2002), 1–14.
- [5] S. Gudmundsson and E. Kappos, On the geometry of tangent bundle with the Cheeger-Gromoll metric, *Tokyo J. Math.*, 25 (2002), 75–83.
- [6] O. Kowalski Curvature of the induced Riemannian metric on the tangent bundle of a Riemannian manifold, *J. Reine Angew. Math.*, 250 (1971) 124–129.

- [7] C. Oniciuc, The tangent bundle and harmonicity, *An. Stiint. Univ. Al. I. Cuza Mat.*, 43(1997), 151–172.
- [8] P. Petersen, *Riemannian geometry*, Springer-Verlag, 1998.
- [9] S. T. Yau, Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry, *Indiana M. J.* 25 (1976), 659–670.

DÉPARTEMENT DE MATHÉMATIQUES, UMR 6205, UNIVERSITÉ DE BRETAGNE OCCIDENTALE, 6, AVENUE VICTOR LE GORGEU, CS 93837, 29238 BREST CEDEX 3, FRANCE

E-mail address: Michele.Benyounes@univ-brest.fr and Eric.Loubeau@univ-brest.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, HESLINGTON, YORK YO10 5DD, U.K.

E-mail address: cmw4@york.ac.uk